

5. Louie D , Earp B & Philip Blazar P Long-term outcomes of carpal tunnel release: a critical review of the literature *HAND* (2012) 7:242–246
6. Marshall S, Tardif G, Ashworth N. Local corticosteroid injection for carpal tunnel syndrome. *Cochrane Database Syst Rev*. 2007(2):CD001554.
7. Page MJ, Massy-Westropp N, O'Connor D, Pitt V. Splinting for carpal tunnel syndrome. *Cochrane Database Syst Rev*. 2012(7):CD010003.
8. Shi Q, MacDermid JC. Is surgical intervention more effective than non-surgical treatment for carpal tunnel syndrome? A systematic review. *J Orthop Surg Res*. 2011;6:17.
9. Stark H, Amirfeyz R. Cochrane corner: local corticosteroid injection for carpal tunnel syndrome. *J Hand Surg Eur Vol*. 2013;38(8):911-4.
10. Royal College of Surgeons:
<https://publishing.rcseng.ac.uk/doi/10.1308/rcsbull.2017.28>
11. Verdugo RJ, Salinas RA, Castillo JL, Cea JG. Surgical versus non-surgical treatment for carpal tunnel syndrome. *Cochrane Database Syst Rev*. 2008(4):CD001552.

N. Dupuytren's contracture release in adults

Updated description of the intervention

NICE recommends no treatment is necessary for people with Dupuytren's disease who do not have contracture. Referral to hand surgery should be made for people with Dupuytren's contractures according to the criteria listed below.

Updated clinical criteria

Summary of intervention

Dupuytren's contracture is caused by fibrous bands in the palm of the hand which draw the finger(s) (and sometimes the thumb) into the palm and prevent them from straightening fully. If not treated the finger(s) may bend so far into the palm that they cannot be straightened. All treatments aim to straighten the finger(s) to restore and retain hand function for the rest of the patient's life. However none cure the condition which can recur after any intervention so that further interventions are required.

Splinting and radiotherapy have not been shown be effective treatments of established Dupuytren's contractures.

Several treatments are available: collagenase injections, needle fasciotomy, fasciectomy and dermofasciectomy. None is entirely satisfactory with some having slower recovery periods, higher complication rates or higher reoperation rates (for recurrence) than others. The need for, and choice of, intervention should be made on an individual basis and should be a shared decision between the patient and a practitioner with expertise in the various treatments of Dupuytren's contractures.

No-one knows which interventions are best for restoring and maintaining hand function throughout the rest of the patient's life, and which are the cheapest and

most cost-effective in the long term. Ongoing and planned National Institute for Health Research studies aim to address these questions..

Number of CCG interventions in 2017/18

14,376

Recommendation

- Treatment is not indicated in cases where there is no contracture, and in patients with a mild (less than 20°) contractures, or one which is not progressing and does not impair function.
- An intervention (collagenase injections, needle fasciotomy, fasciectomy and dermofasciectomy) should be considered for:
 - a. finger contractures causing loss of finger extension of 30° or more at the metacarpophalangeal joint or 20° at the proximal interphalangeal joint.
 - or**
 - b. severe thumb contractures which interfere with function
- NICE concluded that collagenase should only be used for:
 - a. Participants in the ongoing clinical trial (HTA-15/102/04)
 - or**
 - b. Adult patients with a palpable cord if:
 - i. there is evidence of moderate disease (functional problems and metacarpophalangeal joint contracture of 30° to 60° and proximal interphalangeal joint contracture of less than 30° or first web contracture) plus up to two affected joints;
 - and**
 - ii. needle fasciotomy is not considered appropriate, but limited fasciectomy is considered appropriate by the treating hand surgeon

Rationale for recommendation

Contractures left untreated usually progress and often fail to straighten fully with any treatment if allowed to progress too far. Complications causing loss, rather than improvement, in hand function occur more commonly after larger interventions, but larger interventions carry a lower risk of need for further surgery.

Common complications after collagenase injection are normally transient and include skin breaks and localised pain. Tendon injury is possible but very rare. Significant complications with lasting impact after needle fasciotomy are very unusual (about 1%) and include nerve injury. Such complications after fasciectomy are more common (about 4%) and include infection, numbness and stiffness.

References

1. http://www.bssh.ac.uk/_userfiles/pages/files/Patients/Conditions/Elective/dupuytren's_disease_leaflet_2016.pdf
2. <https://cks.nice.org.uk/dupuytren's-disease>
3. Crean SM, Gerber RA, Le Graverand MP, Boyd DM, Cappelleri JC. The efficacy and safety of fasciectomy and fasciotomy for Dupuytren's contracture in European patients: a structured review of published studies. *J Hand Surg Eur Vol.* 2011;36(5):396-407.
4. Krefter C, Marks M, Hensler S, Herren DB, Calcagni M. Complications after treating dupuytren's disease. A systematic literature review. *Hand surgery & rehabilitation.* 2017, 36: 322-9.
5. NICE 2004. Needle fasciotomy for Dupuytren's contracture. <https://www.nice.org.uk/guidance/ipg43>
6. NICE, 2017. Collagenase clostridium histolyticum for treating Dupuytren's contracture. : <https://www.nice.org.uk/guidance/ta459>,
7. Rodrigues JN, Becker GW, Ball C, Zhang W, Giele H, Hobby J, et al. Surgery for Dupuytren's contracture of the fingers. *Cochrane Database Syst Rev.* 2015(12):CD010143.
8. Scherman P, Jenmalm P, Dahlin LB. Three-year recurrence of Dupuytren's contracture after needle fasciotomy and collagenase injection: a two-centre randomized controlled trial. *J Hand Surg Eur Vol.* 2018;43(8):836-40.
9. Skov ST, Bisgaard T, Sondergaard P, Lange J. Injectable Collagenase Versus Percutaneous Needle Fasciotomy for Dupuytren Contracture in Proximal Interphalangeal Joints: A Randomized Controlled Trial. *J Hand Surg Am.* 2017;42(5):321-8 e3.
10. Stromberg J, Ibsen Sorensen A, Friden J. Percutaneous Needle Fasciotomy Versus Collagenase Treatment for Dupuytren Contracture: A Randomized Controlled Trial with a Two-Year Follow-up. *J Bone Joint Surg Am.* 2018;100(13):1079-86.
11. van Rijssen AL, Gerbrandy FS, Ter Linden H, Klip H, Werker PM. A comparison of the direct outcomes of percutaneous needle fasciotomy and limited fasciectomy for Dupuytren's disease: A 6-week follow-up study. *J Hand Surg Am.* 2006, 31: 717-25.
12. van Rijssen AL, ter Linden H, Werker PM. Five-year results of a randomized clinical trial on treatment in Dupuytren's disease: Percutaneous needle fasciotomy versus limited fasciectomy. *Plast Reconstr Surg.* 2012, 129: 469-77.

O. Ganglion excision

Updated description of the intervention

Most people live comfortably with ganglia and they often resolve spontaneously over time. Ganglion excision can be unnecessary, can cause complications, and recurrence is common following surgery. The complications may be similar to or worse than the original problem. Ganglion excision should only be offered under the criteria outlined below.

Updated clinical criteria